

Review Article

# Paclobutrazol Boost up for Fruit Production: A review

Ashok Kumar<sup>1\*</sup>, Sant Ram<sup>2</sup>, LD Bist<sup>2</sup> and CP Singh<sup>2</sup>

<sup>1</sup>Professor Horticulture, Department of Agriculture, DBIMS, Manduwala, Dehradun-248007, Uttarakhand, India

<sup>2</sup>Professor Horticulture, College of Agriculture, G.B.P.U.A &T- Pant Nagar, U.P, India

\***Address for Correspondence:** Ashok Kumar, Professor Horticulture, Department of Agriculture, DBIMS, Manduwala, Dehradun-248007, Uttarakhand, India, Tel: +91-8983232585; Email: yadavakdr@gmail.com

**Received:** 27 August 2020; **Accepted:** 28 October 2020; **Published:** 18 December 2020

**Citation of this article:** Kumar A, Ram S, Bist L, Singh C (2020) Paclobutrazol boost up for fruit production: A review. Rea Int J of Energy Environ Sci. 1(1): 019-031. DOI: 10.37179/rijees.000005.

**Copyright:** © 2020 Kumar A, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

Paclobutrazol (PBZ), a triazole derivative, has been effectively used to induce and manipulate flowering, fruiting, and tree vigour in several perennial fruit crops. Soil application of paclobutrazol has been efficacious in promoting flowering and increasing yield in many fruit crops. However, there are some conflicting reports on its impact on fruit quality parameters. Besides reducing gibberellins level, PBZ increases cytokinin contents, root activity and C: N ratio, whereas its influence on nutrient uptake lacks consistency. The ability of the crops to produce fruits throughout year is of great interest in recent years. Usually it is applied as a soil application in the month of September-November in case of mango. It inhibits gibberellins biosynthesis at kaurene stage and has proved to be reduction of vegetative growth, promising for flower initiation in shoot bud, giving early and profuse flowering, increases fruit yield and improving quality regularly in alternate bearing cultivars. The main aim of this review is to focus upon contemporary information about paclobutrazol in fruit production. The persistence of cultar in orchard soil for a long time and its half-life varies with soil type and climatic conditions, which may severely affect the development of subsequent crops and moved rapidly between the shoot tips and basal nodes both in the acropetal and basipetal directions and persisted for several months. The potential of PBZ to contaminate groundwater at optimum concentrations is low however the risk of its exposure to aquatic life is high. PBZ is considered moderately hazardous for human beings with remote chance of being genotoxic and carcinogenic. In view of the above, optimized use of the PBZ to derive maximum benefit with least undesirable impact on food and environmental safety aspects is suggested.

**Keywords:** Growth retardant, water relations, nutrient uptake, fruits, PBZ, Cultar, paclobutrazol; PP333; vegetative growth; fruit production

## Introduction

Plant growth retardants are being used widely in chemical manipulation of growth and development by modifying associated biochemical and physiological processes. Among them, paclobutrazol is considered as one of the most versatile plant growth retardant which restricts vegetative growth and induce flowering in many fruit crops like apple and pear [1], peach [2], citrus [3] and mango [4]. It restricts induced tree vigour and flowering responses which have been reported as the consequences of modifications in physiological activities as well as changes in cellular metabolites [5]. Mango (*Mangifera indica* L.) belonging to the family Anacardiaceous, is the

most important commercial fruit of India and considered as King of fruit, because of its rich, luscious, aromatic flavor and a delicious taste in which sweetness and acidity are delightfully blended. It is the most popular and the choicest fruit and occupies a prominent place among the fruits of the world. In India it is grown on an area of 2.516 million hectares with annual production of 18.431 million tone having productivity of 7.3 metric tons per hectare Anonymous [6]. Although, alternate bearing is a major problems in mango production and its means "a condition at which high or optimum fruit production in on year or higher and certain year bear little or no fruit (off year), but growth regulators such as cultar reported to be effective on inducing flowering mango off year (Sinde, et al. 2000).

The alternate flowering in mango might be due to improper orchard management practices, environmental factors, varietal character, or imbalance of hormone, either alone or in combination. The ability to produce crops throughout the entire year is of great interest for mango production under sub-tropical and semi-arid conditions. The biennial bearing is profoundly serious problem in north, east and central Indian commercial cultivars, while most of the south Indian varieties bear regularly. Therefore, application of cultar is most widely studied in view of its high potential for controlling plant growth and development of fruit crops in general and mango production. It is applied either in the soil or as foliar spray in the September-November.

The persistence of cultar in orchard soil for a long time and its half-life varies with soil type and climatic conditions, which may severely affect the development of subsequent crops and moved rapidly between the shoot tips and basal nodes both in the acropetal and basipetal directions and persisted for several months. It inhibits gibberellin biosynthesis at kaurene stage and has proved to be reduction of vegetative growth, promising for flower initiation in shoot bud, giving early and profuse flowering, increases fruit yield and improving quality regularly in alternate bearing cultivars. The main aim of this review is to focus upon contemporary information about cultar in mango production. There are several strategies that control the problems of flowering periodicity and tree vigor in tropical and sub-tropical region such as the use of shoot pruning practices, dwarfing rootstock and growth regulators, the use of plant bio regulator is the most promising approach for managing canopy and ensuring regularity in flowering and enhancing fruit yield under commercial cultivation (Olivier, et al. 1990). Among the cellular metabolites, accumulation of phenols in vegetative organs and altered biochemical balance are important in restriction of vigour in mango [7] and induction of flowering [8].

Paclobutrazol (PBZ) is a triazole derivative with the empirical formula [(2RS, 3Rs) -1 - (4- chlorophenyl) 4,4-dimethyl-2- (1H-1, 2, 4-triazole-1-yl) pentan-3-ol], which plays an important role in regulating excessive vegetative growth, enhancing and advancing flowering, inducing early bearing, managing biennial bearing tendency, establishing a high density plantation. The application of paclobutrazol to soil promotes flowering and increasing yield in many fruit crops. Besides reducing gibberellins level, paclobutrazol increases cytokinin contents, root activity and C: N ratio, whereas its influence on nutrient uptake lacks consistency. It also affects microbial population and dehydrogenase activity in soil. PBZ has been characterized as an environmentally stable compound in soil and water environments with a half-life of more than a year under both aerobic and anaerobic conditions. However, when it is applied in optimized rate the residual concentration detected will not be above quantifiable level (0.01 ppm) in soils and fruits. Cultar is effective not only in flower induction but also in early and off-season flower induction in mango [9-13]. However, the Mode of action of plant growth regulators such as cultar, is highly specific to cultivar, rate of application, cultivar, developmental stages, and climatic condition (Hoffmann 1992).

Thus, cultar holds considerable promise in manipulation of flowering, yield and vigour in fruit crops. However, its high potency for harmful to nature, slow mobility in the orchard soil, persistence in

soil and fruit over its long-term use (USEPA 2007). Hence, an effort was made to review the research work on the use of cultar in mango production. Physical and chemical properties of cultar Cultar is a plant growth inhibitor belonging to the triazole group.

### Application Methods of Cultar

There are four application methods of cultar such as soil application, foliar application, trunk application, injection methods, out of which soil and foliar application mostly use in mango orchard. The application of cultar to soil as a drench around the tree trunk (TSLP) is the most effective method, as it ensures proper uptake by the tree. The required quantity is mixed in approximately one litre of water and poured onto the soil around the trunk in a circular band. It is a triazole derivative being capable to inhibit the biosynthesis of gibberellins potentially (Sinde, et al. 2000) and has been effectively used in reducing canopy volume and increasing flower intensity in mango ([14]. Cultar is effective not only in flower induction but also in early and off-season flower induction in mango [9-14]. However, the Mode of action of plant growth regulators such as cultar, is highly specific to cultivar, rate of application, cultivar, developmental stages, and climatic condition (Hoffmann1992).

Thus, cultar holds considerable promise in manipulation of flowering, yield and vigour in fruit crops. However, its high potency for harmful to nature, slow mobility in the orchard soil, persistence in soil and fruit over its long-term use (USEPA 2007). Hence, an effort was made to review the research work on the use of cultar in mango production. Physical and chemical properties of cultar Cultar is a plant growth inhibitor belonging to the triazole group. Fonseca [15] reported that the effects of cultar (0.5, 1.0, 1.5 and 2.0 ml), applied either in the soil or as foliar spray, on the flowering and yield of mango cv. Tommy Atkins Therefore, soil application by trunk soil line pour (TSLP) (Tukey, 1983; Ferree and Schmid, 1988; Kim., et al. [16] and soil drench [17] methods were attempted.

### The Mode of Action

Paclobutrazol inhibits gibberellins biosynthesis by blocking the conversion of kaurene and kaurenoic acid,

**Figure: Mango (Fruit Growth and Quality)**



which inhibits cell elongation and internodes extension and ultimately retards plant growth. Gibberellins stimulate cell elongation. When gibberellins production is inhibited, cell division still occurs, but the new cells do not elongate. That result in the production of shoots with the same numbers of leaves and internodes compressed into a shorter length. Even reduction in the diameter of the trunk is noticed. Paclobutrazol treated trees shows increased production of the hormone abscisic acid and the chlorophyll component phytol, which are beneficial to tree growth and health. It also induce morphological modifications of leaves, such as smaller stomatal pores, increased number and size of surface appendages, thicker leaves, and increased root density that may provide improved environmental stress tolerance and disease resistance and it also has some fungicidal activity due to its capacity as a triazole to inhibit sterol biosynthesis [18].

### Translocation of PBZ in plant

PBZ is applied as a soil drench (application to roots, more popular and convenient) through trunk injection (directly to the vascular system of the stem using pressure). Through xylem it translocates to other parts of plant, however a few research evidences have been provided to support this assumption. Gas chromatography-mass spectrometry confirmed that PBZ was taken up by roots and transported primarily through xylem to stems and accumulated in leaves.

Influences the mechanism of nutrient uptake The ability of roots to draw nutrients from the soil and to deliver these to the aerial plant tissues at a rate that matches the needs of growth is key to ensure physiological growth and development of plant, Whereas mismatch between the demand of the shoot and the supply from the roots can affect productivity (Tester and Leigh 2001). Kotur [19] observed significant increase in the root activity towards the trunk and close to soil surface and sparser root activity in the subsoil zone and in drip line area in paclobutrazol treated mango plants. On the other hand, Werner [20] observed an increase of N, Ca, Mn, Zn and B contents and decrease of P, K and Cu contents in cultar treated mango trees. Soil application of cultar (2.0-8.0 g ai) for two consecutive years in mango increased the levels of phosphorus, potassium and calcium at lower doses but decreased at higher dose. The findings indicate inhibitory effect of paclobutrazol at higher concentration on soil nutrient status and microbial population [21].

PBZ also promotes the avoidance of salt stress in mango by increasing the levels of photosynthetic pigments, water content, K<sup>+</sup> uptake and uptake of harmful Na<sup>+</sup> and Cl<sup>-</sup> ions (Kishor, et

al. 2009). The influence of paclobutrazol on leaf nutrient content lacks consistency as it showed variation with the crop species and soil conditions. Alter the phytohormone/ endogenous hormone Phytohormones play a crucial role in regulation of plant growth and development. There is an increasing evidence for a decisive function of certain hormones in the establishment of developmental programs of plants. Gibberellins are destined for vegetative growth, whereas cytokinin induces reproductive phase (Alabadi, et al. 2009). The relative concentration of gibberellin and cytokinin decides the fate of the shoot. A significant decline in the GA3-like compounds was observed in the shoots of PBZ-treated plants after two months of application in mango and there was no difference in the level of GA3-like substance between control and treated plants one year after the treatment.


This suggests the need for repeat application of cultar [9]. Upreti, et al. [22], reported that cultar besides affecting gibberellins also increases ABA and cytokinin, viz. zeatin (Z), zeatin riboside (ZR) and dehydrogenation riboside (DHZR), contents concomitant with C: N ratio and leaf water potential in mango buds to elicit flowering responses. In a similar findings, Singh and Sharma (2008) recorded increase in C:N ratio, leaf water potential, chlorophyll content, total sugar, total protein, nitrate reductase activity, ABA and cytokinin's - zeatin (Z), zeatin riboside (ZR) and dehydrogenation riboside (DHZR) in paclobutrazol treated mango. Adil, et al. (2011) also recorded enhancement in the levels of zeatin (z), zeatin riboside (zr), isopentenyl Adenosine (i-Ado), isopentenyl Adenine (i- Ade), and abscisic acid (ABA), through at low level, along with the increase in starch and sugar contents in cultar treated trees of mango during the floral induction period. Whereas gibberellins (GA1+3+20) and auxin (IAA) were decreased during the same period. Cultar also induces morphological modifications such as enhanced leaf specific weight, stomatal density, leaf thickness, root-to-shoot ratio and root density that strengthen stress tolerance capacity in plants. Additionally, it has also fungicidal activity due to its inhibition of sterol biosynthesis [18], Fernandez, et al. 2006).

### Effects of PBZ on various tree attributes

In tropical fruit orchards, it is desirable to control the vegetative growth and to reduce the canopy size since small trees capture and convert the sunlight into fruit biomass in a better way than larger trees because of more surface area. Increase in production with enhanced fruit quality can be achieved by managing the tree canopy. Manipulation in tree physiology with the use chemical growth retardants has been considered as an important determinant of productivity enhancement in many fruit crops. Application of paclobutrazol in the soil has been commercialized for early and enhanced flowering in some of the fruit crops.

### Plant growth and vigour

Paclobutrazol prevents stem elongation [23]. PP333 decreased the elongation of new shoots in peach and the effect increased with increasing concentration from 500 to 1500 ppm Chen, et al. [24,25], also noted that foliar sprays of PP333 at 1000 or 2000 mg a.i. per litre suppressed peach seedling growth. Allan, et al. [26], also observed significant reduction in competitive early vegetative growth by a soil drench of PP333 in peach cv. Flordaprince. Similarly, Biasi, et al. [27] and George, et al. [28] observed reduction in growth following



**Figure 1:** [Paclobutrazol (PBZ), [2RS, 3RS]-1-[4-chlorophenyl]-4, 4-dimethyl-2-(1H-1, 2, 4-triazol-1-yl) pentan-3-ol].



**Figure 2:** Paclobutrazol (PBZ) packaging in dust Form.

PP333 treatment in peach and nectarine, respectively. Lever, et al. [29] found more than 50 per cent reduction in shoot growth of Red Delicious apple following PP333 foliar spray at the rate of 750 ppm. However, the effectiveness of PP333 varied with the dose, time, and method of application. Irving and Pallesen [30], found that on two-year-old apple 1000 ppm of PP333 had extraordinarily little effect on vegetative growth and remained effective up to 82 days of application. But Stinchcombe, et al. [31], reported that, in Cider apple, PP333 at 2000 ppm remained effective in the following year also. ElKhoreiby, et al. [32], recorded maximum retardation in growth when PP333 was applied 21 days after petal fall.

The reduction in shoot length of Fuji apples was also reported by Kim, et al. [16]. Greene [33], observed the growth retardation effect of PP333 continuously for three years on apple trees when it was applied at higher rates of 1500-3000 ppm as foliar sprays. Mavrodiiev and Manolov [34], reported that PP333 was less effective in controlling growth during the year of application but more effective during the following year. Quinlan and Richardson [35], found that 14C PP333 translocated acropetally when applied to young stem internodes and to a lesser degree, from the youngest unrolled leaf, however, there was no label moved from mature leaves. Paclobutrazol is also reported to regulate the vegetative growth of peach and cherry [36,37]. Sharma and Joolka [38], and Sharma and Joolka [39], recorded reduced extension growth, plant height and plant spread with paclobutrazol in Non Pareil almond plants. Mir, et al. [40], reported that paclobutrazol significantly retarded the shoot growth, shoot diameter and trunk cross-sectional of 'Roundel' apricot trees growing under low density planting system.

### Effect on vegetative growth

Many investigations have revealed the beneficial effects of PBZ in restricting vegetative growth and successful induction of flowering in apple, mango, grape etc. The application of paclobutrazol (1500 to 3000 ppm) at full bloom and 21 days after full bloom resulted in the reduction of shoot growth in 'Golden Delicious' apple [41]. Quinlan and Richardson [42], inferred that application of paclobutrazol at 500 ppm alone was effective in reducing the shoot length (9.5 cm) and the combination with GA3 was not effective in apple seedlings. Five-year-old MM.106 (*Malus domestica* Borkh.) trees growing under a high-

density (10000 trees/ha) planting system treated with paclobutrazol at 250 mg per tree in August by Khurshid, et al. 1997 [43], showed reduced number of total shoots and buds. This showed that PBZ can be used to manipulate apple tree growth in a high-density apple production system. Paclobutrazol when applied during early summer has been observed as an effective suppressant of stem growth in sweet cherry [44]. Similarly, Webster, et al. [35], reported that application of paclobutrazol at 1.6 g a.i. tree<sup>-1</sup> and followed by 0.8 g a.i. in next year inhibited extension of growth in young cherry trees on either colt or FB22 rootstocks. 2-year-old nashi trees treated with paclobutrazol as soil drench and foliar sprays [45]. The cultivars treated were 'Hosui', 'Kosui', 'Nijisseiki', and 'Shinsui'. All cultivars showed a significant reduction in vegetative growth within the first season and for up to 4 years after initial application. Most reduction in growth was obtained from soil applications. Least reduction in growth was from a foliar application at the lower rate of 125ppm.

Application of paclobutrazol on 'Redhaven' cultivar of peach reduced terminal growth and advanced leaf fall [46]. Similarly, the vigour of mango was consistently reduced with paclobutrazol application in a range of Indian cultivars [47]. The soil drenching with paclobutrazol at the rates of 12, 10, and 8 g a.i. suppressed the vegetative growth, canopy volume, and flush length of reproductive shoots, fruit setting, panicle length as compared to control in mango [12]. Similar result was observed by Teferi, et al. (2010) in Tommy Atkins mango with maximum effect at 8.25 g a.i. per tree. Soil application of paclobutrazol recorded significant reduction in canopy volume by noticeable reduction in number of shoots per terminal and also checked the growth of new shoots [48]. Similarly, the growth inhibitory response of PBZ reported in different varieties of mango [4], could be the consequences of modification in photosynthesis rate [49] and carbohydrates [5], besides reductions in gibberellins [22].

Leaf area: Biasi, et al. [27], applied 0.1 g PP333 per seedling as soil drench to Nemaguard peach and noted reduction in leaf area. Curry and Williams [50], found reduced leaf size of Well Spur Delicious apple with the application of 20 g PP333 per 9.5 m<sup>2</sup> as soil drench. Similar observations in apple have been made by Stinchcombe, et al. [31], Swietlik and Miller, Chogtu Greene, Abod and Webster, Bhatia and Xia, et al. (1994), Val, et al. (1999) [33, 51-54], reported 40 per cent reduction in leaf area and 29 per cent reduction in dry weight of peach with paclobutrazol application. Root growth: Soil drenching with 0.1 g of PP333 per plant resulted in reduced fresh and dry weight of root in containerized peach cv. Nemaguard seedlings [27]. Contrary to this, promotion in the root growth of apple trees with PP333 has been reported by Lenz [55], Steffens, et al. [56] and Lehman, et al. [57]. ElHodairi, et al. [58], found an increased root: shoot ratio and this was associated with a redistribution of 14C-labelled assimilates in the plant. Dry matter accumulation in the roots, particularly in lateral and fibrous roots, increased with 1.5 g a.i. and 0.75 g a.i. PP333 applied in spring as soil drench in Aki Fuji apples [59]. Similarly, Curry and Williams [60], recorded an increased root dry weight with the lower dose of PP333 but not with its higher dose.

Whereas Swietlik and Miller [61], reported that total root surface of apple seedling was not affected by PP333. Similarly, Abod and Webster [53], found no effect of 500 ppm PP333 spray on the root weight of MM106 apple rootstock. Bhatia [54], recorded reduced root length in MM109 apple rootstock treated with 1.0 g PP333 as

soil drench. Similarly, Zeller, et al. [62], observed decrease in root growth of potted plants of Smoothie Golden Delicious apple when treated with 1, 10 or 100 mg PP333 per plant. Sharma and Joolka [38], recorded reduced total root length and root dry weight with paclobutrazol in Non Pareil almond plants.

### Flower Bud Differentiation

Flowering in mango is preceded by the differentiation of the flower bud in the shoots. Physiological maturity appears causally related with flower bud differentiation (Muhammad, et al. 1999). In Indian sub-continent, the time of the flower bud differentiation has been reported by various workers to varying from October to December. In Baramasi mango bud differentiation, most often, takes place twice a year, i.e. during May- June and September-October. Fluctuation in the time of flower bud differentiation is dependent on genetic, environmental, and endogenous tree factors (Chaco, 1991; Schaffer, et al. 1994). Thus, flower bud differentiation seems to be depending on the fluctuations in temperatures and the fruit load borne by the tree during previous year.

Ravishankar, et al. (1979) found that flower bud differentiation in Alphonso mango, a irregular or erratic bearing cultivar grown under the mild tropical climate of Dharwar (India) was initiated in early October and reached a peak by November. Singh [63], reported the last week of December to be the critical time for flower bud differentiation under north Indian climate. Regulation of flowering in off season the application of cultar before flower bud differentiation or three months earlier than anticipated flowering has been effective in inducing flowering in mango without accompanying reduction in shoot length. However higher concentration leads to canopy and panicle compaction [64].

Chusri, et al. [66], reported that in the PBZ treated trees bud break occurred 18-22 days earlier than in the control trees in 'Irwin' mango. Apical bud breaking exceeded 98% in the PBZ-treated trees compared

with only 42% in the control trees. More than 96% of the PBZ-treated trees produced floral shoots, compared with only 35% of the control trees. The panicles of the PBZ- treated trees were shorter than those of the control trees. Apart from enhancing flowering intensity, cultar has also been effective in increasing sex ratio, cauliflory and axillary flowering in mango (Singh 2000) [66]. Cardoso [67], observed that there was a higher percentage of flowering and fruit production as compared with the control. The PBZ treatment also anticipated flower initiation and fruit harvest, which means that out of season production, provides higher prices and more profitability.

Reddy and Kurian [68], observed that under tropical climate, application of paclobutrazol for three consecutive years and then its discontinuation for the subsequent three years appears in twenty years old mango trees to be appropriate. However, the continuous optimum use of cultar in high density planting is imperative to manage canopy and to induce precocious flowering as it was also observed that young plants respond better than old ones. Soil application around the tree trunk (collar drench) was more efficacious than foliar application as it ensures proper uptake in inducing flowering and fruiting (Kulkarni, et al. 2006). On the other hand, Yeshitela [11], reported that application of cultar both as a soil drench and foliar application were effective in suppressing vegetative growth and enhancing flowering, yield, fruit quality as well as number of hermaphrodite flowers in mango.

The response to cultar varied with cultivar and crop load. The shoot retarding effect of cultar was generally limited in mango var. Sensation but was pronounced in Tommy Atkin. Moreover the average fruit weight and yield were increased with the rate of paclobutrazol in Sensation, whereas fruit weight and yield were reduced in Tommy Atkin (Singh and Bhattacharjee 2005). Singh, et al. [21], reported soil application of PBZ at 5 and 10 g/tree considerably increased the percentage of panicles and hermaphrodite flowers on twenty-five-year-old Dashehari mango trees. Soil application of paclobutrazol at 5 g/tree was most effective to induce a greater number of flowering shoots in mango cv. Gulab Khas 69].

Mouco, et al. [70], observed that Paclobutrazol promoted flowering in mango trees in any season of the year, under tropical semi-arid conditions, but its efficiency was related to the maximum and minimum air temperatures at the time of bud break. Bagel, et al. [71], observed maximum number of flowering shoots/m<sup>2</sup> (n=30.32) and percentage of flowering shoots (96.15%) when Cultar applied alone (2.50, 3.75 and 5.00 g/ha) and in combination with NAA (20 ppm), on the flowering and fruiting of 10-year-old mango cv. Langra trees and they recorded flowering and fruiting was significantly increased by 7- 30%. Soil application of Cultar promoted flowering, along with cauliflory and axillary flowering [66]. Four months later and monthly thereafter, 10 shoots were dipped in 2% potassium nitrate to assess flowering response. Flowering started only by November or 6 months after paclobutrazol application. By December, all trees treated with 1-gram paclobutrazol per canopy diameter flowered in contrast to only 38% of the control trees. Starch content increased in stems of paclobutrazol-treated trees, suggesting that paclobutrazol promotes flowering by increasing starch accumulation [9]. The panicle size was reduced with the application of Cultar whereas the percentage of hermaphrodite flowers, fruit set and yield was increased.

### PBZ effect on flowering parameters and yield

The fruit set was increased in paclobutrazol treated trees @ 1500



**Figure 3.4:** [Paclobutrazol (PBZ), [2RS, 3RS]-1-[4-chlorophenyl]-4, 4-dimethyl-2-(1H-1, 2, 4-triazol-1-yl) pentan-3-ol]

and 3000 ppm due to an increase in initial fruit set in delicious apple [33], similar results was observed by Elfying, et al. [72], in McIntosh apples that the foliar application of paclobutrazol reduced pre harvest drop when applied within 5 weeks after full bloom. Stan, et al. [74], reported that foliar and soil application of paclobutrazol enhanced the flower bud formation and fruit set-in high-density planting of sweet cherry. In avocado, paclobutrazol enhanced the fruit set by increasing the portioning of dry matter to fruits [74].

Jindal and Chandel [75], applied paclobutrazol in 'Santa Rosa' plum at 125, 250 and 500 ppm once at full bloom and again at pit hardening stage and reported maximum fruit weight of 24.33 g and fruit volume of 21.6 cc in fruits treated with 500 ppm paclobutrazol. Ratna and Bist [76], reported that application of 0.15 g a.i. paclobutrazol cm<sup>-1</sup> trunk diameter increased fruit yield of 'Gala' pear and during the next year, yield was significantly increased with the same application. They also noticed that paclobutrazol 0.3 g a.i. cm<sup>-1</sup> trunk diameter increased the yield by more than 1.35 times during both the years. Arzani, et al. [77] reported that paclobutrazol application advanced flowering of five-year-old vigorous 'Sundrop' apricot trees by 2-4 days and also increased the fruit set, final fruit number, crop density and yield efficiency. Selva strawberry cultivar using paclobutrazol (0,100 mg l<sup>-1</sup>) and other nutrient combination indicated that vegetative growth was reduced with application of paclobutrazol and highest vitamin C was obtained at concentration of 0-100 mg l<sup>-1</sup> PP333 [78].

Kulkarni [47], observed that there was a significant increase in yield of mango per tree by the soil application of paclobutrazol (10 g a.i./tree). In terms of fruit size and quality for at least two years in five years old bearing trees. Effect of PBZ on promotion of flowering in citrus was studied by Fuentes, et al. [79] and result revealed that PBZ significantly increased the percentage of sprouted buds and leafless floral shoots (both single flowered shoots and inflorescence) and reduced the number of vegetative shoots. The application of paclobutrazol at 1 g a.i./m of canopy diameter increased the female inflorescence production (18.10%) without negative effect on fruit set (90.68%) in 'Eviarc Sweet' cv. of jackfruit. Female inflorescences were produced in the off-season (August and September) which was not observed in untreated trees.

[80]. Among the chemicals suggested, paclobutrazol is considered as one of the most versatile plant growth retardant which restricted vegetative growth and induced flowering in many fruit crops like apple and pear [1], peach [2], citrus [3] and mango [4]. Early and intense flowering induced by PBZ may be the consequence of early shoot maturity and increased photosynthesis rate [81], carbohydrate accumulation [82] and decline in flowering reducing hormone, gibberellins [22], profuse and early flowering [10,11,14,47,68,83-87], reduced panicle length [4, 12, 88, 89], increased the number of perfect flowers and fruit set [66,89-91], were reported in various fruit crops.

All the available evidence opined that carbohydrate reserves played an important role in flower bud differentiation and they provide conditions favorable for the synthesis of substances which are required for flower bud differentiation [92,93]. The high C: N ratio during flower bud differentiation was ascribed to the increased carbohydrate availability [94] and is considered as an important factor in regulation of flowering in fruit crops [95,96]. Paclobutrazol is known to decrease vegetative growth rate through early cessation

of growth, which results in the accumulation of carbohydrates in trees and slightly decreasing the total nitrogen in the terminal shoots, which favours flowering by maintaining high C: N ratio in the shoots.

The C: N ratio differs with growth of shoots in the varieties revealing its dependence on environmental conditions and prevailing metabolic balance. The paclobutrazol induced enhancement in C: N ratio has been reported in mango [5,9,22,82, 97,98] and in pummelo [99], distinct differences in carbohydrate pattern are seen in vegetative growing shoots and flowering shoots. Shoots that are going to differentiate into flower buds are the growing sinks and the actively dividing cells of induced flower buds require high energy [100]. Apparent increase in sugar levels during floral induction period has been reported in mango by several researchers [95,96,101]. Consistently higher production of total sugars and reducing sugars with peak availability at bud burst in apical buds of paclobutrazol treated trees is reported in mango [5,104].

Paclobutrazol induced increase in soluble sugars at flowering has also been reported in mango [82]. Among the cellular metabolites, accumulation of phenols in vegetative organs has been depicted as one of the important in imparting of vigour restriction effects in mango [103,104] and also for induction in flowering [8]. The possible mechanism by which phenols exert its effects on tree vigour and regulation of flowering in mango are less understood. However, steady increase in phenol content with advancement of flower bud differentiation has been reported in mango by Palanichamy, et al. [96] and Kumar, et al. [105]. The paclobutrazol induced tree vigour restriction and flowering responses have been reported as the consequences of changes in cellular metabolites [22,82]. High phenol content in terminal buds due to paclobutrazol application restricted the vigour and enhanced the flowering has also been reported by Kurian and Iyer [91].

### PBZ effect on fruit quality

Fruit quality improvement with respect to pulp content, TSS, TSS to acid ratio, total sugars and reducing sugars in response to PBZ application can be related to the assimilate partitioning in plant. The greater suppression of vegetative growth causes assimilates demand in unidirectional manner to the developing fruit, resulting in high quality fruits from PBZ treated plants. Application of paclobutrazol @ 0.33, 0.50, 0.66 and 1.32 g a.i. as soil application to 'Flavorest' peach hastened the fruit colour than control [106]. Similarly, application of 500 mg l<sup>-1</sup> paclobutrazol sprayed within 5 weeks after full bloom to 'McIntosh' apples gave high percentage of fruit with acceptable red color at harvest [72].

Singh and Dillon (1992) reported that soil application of PBZ to Dashehari mango recorded higher fruit yield and high TSS: acid ratio compared to foliar application, while fruit weight: stone and pulp: stone ratio did not differ significantly. Vijaylaxmi and Srinivasan [88], in an experiment with 10 years old Alphonso mango trees treated with paclobutrazol (10 ml), KNO<sub>3</sub> (1%), urea (1%), ethrel (200 ppm), NAA (20 ppm) or mepiquat chloride (500 ppm) found that among all the treatments, paclobutrazol (10 ml) resulted in increased ascorbic acid content, total sugars and reducing sugars, TSS, acidity and sugar: acid ratio in harvested fruits. A significant improvement in the fruit quality of cv. Langra in terms of total soluble solids (TSS), total acidity, total chlorophyll, total carotenoids and α-amylase activity due

to paclobutrazol @ 6 g a.i./tree in comparison to control was reported by Singh and Saini [108]. Further they evaluated the efficacy of soil applied paclobutrazol (2, 4, 6 and 8 g a.i./tree) on Langra cultivar of mango for three consecutive years at Lucknow and reported a significant increase in fruit set, fruit retention per panicle and yield per tree due to PBZ @ 6 g a.i./tree. Saxena, et al. [108], reported that paclobutrazol, a flower inducing chemical, enhanced the catalase and peroxidase activities over the untreated control and maximum enhancement was recorded at 8 g a.i. The decreasing trend of protein with paclobutrazol treatment was recorded in adjacent leaves of flower buds. The results implicated the possible role of catalase and peroxidase and other associated biochemical changes in paclobutrazol induced flowering in mango. The soil drenching of paclobutrazol at 3.0 ml m<sup>-1</sup> canopy diameter to the mango cv. Totapuri was done to study the role of carbohydrates in the paclobutrazol induced floral initiation by Upreti, et al. [5].

The results indicated that paclobutrazol induced flowering was accompanied by an increase in starch in leaf concomitant with increased insoluble sugars like sucrose, glucose and fructose in apical buds as well as inhibition in the amylase activity in association with increase in the activities of acid invertase, sucrose phosphate synthase and sucrose synthase in the apical buds. Similarly in CO 2 papaya (dioecious) there was increase in amino acids, total carotenoids, TSS, sugars, ascorbic acid and sugar-acid ratio as compared to control, the response being linear with the increasing concentrations PP333 as soil drench at two levels viz., 25 and 50 mg a.i./plant [6] The improvement in fruit quality parameters such as high edible portion, longer shelf life, higher TSS, increased vitamin C, lower titrable acidity, high dry matter content and high reducing and total sugars with PBZ was reported by Vijayalakshmi and Srinivasan Hoda, et al. Bamini, et al. Sarkar and Rahim and Reddy, et al. [4,87-89,109], in different varieties of mango. An increase in the contents of ascorbic acid and carotenoids which are documented as potential antioxidants with PBZ application has also been reported in mango [87], papaya [110], guava [111]. However, non-significant effect on fruit quality with PBZ application was reported by Tandel and Patel [48] and Upreti, et al. [22].

### Increase fruit set and yield

Bagel, et al. [71], were recorded maximum yield per tree (68.12 kg), yield per hectare (106.25 q/ha), and yield increase over the control (29.85%) of 10 year old mango cv. Langra trees applied with Cultar at 5.00 g/ha in combination with 20 ppm NAA. Soil application of paclobutrazol at 5 g/tree was most effective to improve the fruit set and Int.J. Curr.Microbiol. App.Sci (2018) 7(2): 1552-1562 1558 [112], fruit retention during the off year. The highest yields of 70.50 and 68.70 kg per tree during the off year were recorded under soil application of paclobutrazol at 5 and 10 g/tree, respectively [69]. Increase in fruit set per panicle is due to retardation of plant vigour by growth retardant.


Benjawan [113], reported that PBZ had no significant effect in extending number of days from flower initiation up to full bloom. PBZ also had no significant effect in delaying fruit maturity age but fruit sets were significantly increased with PBZ rates applied. PBZ had a highly significant effect on fruit length but significantly decreased fruit thickness. Fruit yields were significantly increased with PBZ application. The best application rate of PBZ was found

with T< sub>7 (1000 ppm/plant) with an extended flower raceme length of 5 cm. This treatment gave the highest mango edible fruit yield of 48 281.25 kg ha<sup>-1</sup>. Paclobutrazol was applied in mango cvs. Chausa, Dashehari and Langra as soil drench @ 2, 4, 6 and 8 g/tree and recorded maximum fruits set per panicle, fruit number and fruit yield per tree in 4 g/tree treated Dashehari tree whereas 6 g/tree of PP333 was found more effective in cvs. Chausa and Langra [81]. Soil drenching of PP333, 5 or 10 ml at 120, 90 and 60 days before bud break on mango cv. Neelum and recorded the maximum number of fruits (380) and fruit yield per tree (91.65 kg) both in off season and main season (302 fruits and 72.85 kg per tree) with 5 ml of PP333, 90 days before bud break [114].

### Effects on physiological attributes

Effect on leaf water potential ( $\Psi_w$ ) The PBZ induced increase in  $\Psi_w$  is speculated as the result of increased root hydraulic conductivity, reduced transpiration, and increased ABA levels. Increased ABA reduces the transpiration losses by inducing stomatal closure [115]. As ABA is known to induce stomatal closure and is expected to reduce the water loss through transpiration. The increased water levels due increased ABA are expected to induce bud dormancy which could be of relevance to flower bud differentiation in mango [22,82,103].

Improve Fruit Quality Due to cultar application chemical composition of fruits was considerably enhanced in terms of TSS, total sugars, ascorbic acid, and TSS/acid ratio; however, there was depletion in titratable acidity in freshly harvested fruits. These results are in conformity with the reports of Vijayalakshmi and Srinivasan [88] in mango. Fruit quality of mango (TSS and acid content) increases with paclobutrazol application [13]. The effect was more pronounced in fruits that received 2000 or 3000 mg l<sup>-1</sup> than lower doses. These treatments attained better quality as judged from the total soluble solids, total acidity, ascorbic acid content, total chlorophyll, total carotenoids, and amylase and peroxidase activity from harvest to 12 days of storage at ambient conditions. Chemical parameters of fruits [68], such as TSS and acidity were not affected by cultar, but average weight of a fruit was less in the case of cultar treatments. The effects of cultar applications on fruit size in mango cv. Sensation, and on fruit retention in Tommy Atkins, were investigated in the Northern Province of South Africa. One or 10 ml of paclobutrazol (as Cultar) was diluted with water to 100 ml (0.25 or 2.50 g a.i.), and was applied to a 60-cm-diameter ring of soil around the trunk of 2-year-old trees prior to the initiation of postharvest flushing.



**Figure 5:** Plant Growth Regulator used in Mango Cvs. Dashhari, Langra, Chausa and Fazali.

In Sensation, the average fruit weight (final fruit size) and tree revenue increased with increasing rates of paclobutrazol applied. The number of fruits retained, and yields were not affected. In Tommy Atkins, the number of fruits retained, average fruit weight, yield and tree revenue decreased with increasing rates of cultar applied (Oosthuysen, et al. 1997). The treatment also improved the fruit quality in terms of total soluble solids (TSS), total acidity, total chlorophyll, total carotenoids, alpha-amylase, and peroxidase activity. The relation of cultar on the water use in terms of soil moisture content, gas exchange attributes of Dashehari mango was also studied indicated that the cultar has capacity to retain the moisture to some extent and the trees are less sensitive to fluctuation in water supply and may be better able to withstand drought conditions [81].

### **Influences the mechanism of nutrient uptake**

Werner [20] reported that, cultar treated mango trees showed an increase of N, Ca, Mn, Zn and B contents and decreased contents of P, K and Cu. On the other hand, the significant increase in the root activity towards the trunk and close to soil surface and sparser root activity in the subsoil zone and in drip line area in paclobutrazol treated mango plants was observed by Kotur [19]. Paclobutrazol also promotes the avoidance of salt stress in mango by increasing the levels of photosynthetic pigments, water content, K<sup>+</sup> uptake and uptake of harmful Na<sup>+</sup> and Cl<sup>-</sup> ions [116].

### **Carbohydrate Content**

PP333 influences the carbohydrate contents in plants. Pith and xylem starch deposits in Sudanelli peach increased with PP333 [117]. Vance Delicious and Red Spur Delicious apples had more leaf total sugar with 0.5 g PP333 and leaf starch with 1.0 g PP333 per plant applied as soil drench [54]. However, Xia, et al. (1994) recorded increased leaf sugar and starch contents in Fuji and Starking apples on treatment with 1000 ppm Chenghuabao (PP333). In Spartan apples treated with PP333, Steffens, et al. [118], found higher leaf starch and static sugar contents. Whereas in Top Red Delicious plants treated with PP333, Wieland and Wample (1985) observed reduction in reducing sugars with 150 mg, increase in starch with 25 and 50 mg than those with 150 mg dose of PP333. In apple plants PP333 treatments resulted in the depletion of carbohydrates [120] and starch in shoots [121]. However, Sharma and Joolka [122], observed increased leaf total carbohydrates content with paclobutrazol in Non Pareil almond plants. Nutrient uptake PP333 influences the leaf nutrient status of various temperate fruit crops. Nitrogen: PP333 treatment reduced foliar N concentration in Nemaguard [123], Flordaprince [26,124], Flordaprince and Flordagold peach cultivars [125] and Red Spur Delicious and Vance Delicious apples [54]. However, Atikson and Crisp (1983) found increased foliar N concentration in apple plants treated with PP333. But Swietlik and Miller [51], could not find any effect of PP333 on the foliar N levels of Golden Delicious apples. However, Sharma and Joolka (2011) recorded reduced leaf N content with paclobutrazol in Non Pareil almond plants. Phosphorus: PP333 treatment reduced foliar P concentration in Nemaguard, [123], Flordaprince [26,124], Flordaprince and Flordagold peach cultivars [125] and Red Spur Delicious and Vance Delicious apples [54].

Increased foliar P concentration in apple plants treated with PP333 has been reported by Atikson and Crisp (1983) and Curry [120]. However, Sharma and Joolka [39], recorded reduced leaf P

content with paclobutrazol in Non Pareil almond plants. Potassium: PP333 treatment reduced foliar K contents in Nemaguard ) [123] , Flordaprince peach [26,124], stone fruits Red Spur Delicious, Vance Delicious [54] and Ace Delicious apple (Curry, 1988). Contrary to this, Swietlik and Miller [51], observed increase in K uptake with the addition of 0.2 ppm PP333 to a nutrient solution in which 11-month-old apple seedlings were grown. However, Sharma and Joolka [39], recorded reduced leaf K content with paclobutrazol in Non Pareil almond plants. Calcium: Increased concentration of foliar Ca with PP333 treatment was observed in Nemaguard [123], Flordaprince [26,124], Flordaprince and Flordagold peach cultivars [125] and Red Spur Delicious and Vance Delicious apples [54].

Similar observations regarding the increase in foliar Ca concentrations in various apple cultivars were made by Atikson and Crisp (1983), Swietlik and Miller [51], Curry [120] and [121]. Swietlik and Miller [51], further reported that Ca content in Golden Delicious increased in proportion to the increasing doses of PP333. Sharma and Joolka [39] also recorded increased leaf Ca content with paclobutrazol in Non Pareil almond plants. Magnesium: Foliar Mg content has been reported to increase with PP333 treatment in Nemaguard [123], Flordaprince [26,124], Flordaprince and Flordagold peach cultivars [125] , Red Spur Delicious and Vance Delicious apples [54] and apple plants [121]. But Curry [120] found reduced levels of foliar Mg in apple plants treated with PP333. However, Sharma and Joolka [39], also recorded increased leaf Mg content with paclobutrazol in Non Pareil almond plants.

### **Efficacy of paclobutrazol on reduction of tree canopy**

Garcia [126] opined that the efficacy of cultar in terms of shoot growth and production efficiency depends on the time of pruning. Ram, et al. [127] observed reduction in tree height; shoot length, shoot girth and internodal length when paclobutrazol (12 and 16 ml) applied with pruning (4 or 5 m height) of mango cv. Dashehari trees. Singh, et al. [21], reported that paclobutrazol as soil drenched reduced tree height, shoot length, tree spread and panicle size in mango cv Dashehari.

### **Regulation of flowering in off season**

Soil application of paclobutrazol at 5 g/tree was most effective to induce a greater number of flowering shoots in mango cv. Gulab Khas [69]. Similar reports were obtained by Bagel, et al. [71], in 10-year-old mango cv. Langra trees. Soil application of Cultar promoted flowering, along with cauliflower and axillary flowering [66,128-140].

### **Degradation and persistences in orchard soil**

Reddy and Kurian [68], also observed residual influence of PBZ in soil if applied continuously for three consecutive years and suggested discontinuation of application or to taper down its dose. Singh [21], also detected paclobutrazol residue below permissible limit (0.4898-1.0005 µg/g) in the rhizosphere after two years of application. Degradation and persistences in orchard soil Paclobutrazol are characterized by moderate potential of mobility in soil which enables it is applied in soil unlike other growth regulators (Costa, et al. 2012), however its mobility varied with the soil type. Studies conducted in USA indicate that half-lives of paclobutrazol residues ranged from 450-950 days for orchard soils which indicates poor degradation rate



of PBZ. Paclobutrazol showed low soil adsorption coefficient ( $KD=1.3$  to  $23.0$  ml/g), however adsorption appeared to increase with soil organic matter and a decrease in soil pH. Studies conducted in USA revealed that less than 10% of total PBZ applied were detected in soils between the depths of 60-120 cm, whereas the PBZ ketone metabolite was predominately detected in the subsurface soil layers though at insignificant levels. Sharma and Awasthi [21], detected residues of paclobutrazol in the tree basin soil (0- 15 cm) at the end of each season followed by a slight increase in the number of residues with the year of applications [140-155].

Reddy and Kurian [68], also observed residual influence of PBZ in soil if applied continuously for three consecutive years and suggested discontinuation of application or to taper down its dose. Sharma, et al. (2008) could not detect paclobutrazol residues above quantifiable levels (0.01 ppm) either in tree basin surface soils or in the fruits even after more than five years continuous application. However, they further reported that the residues increased to 0.34 ppm with the increase of the application rate (20 g a. i. /tree). Singh and Bhattacharjee (2005) also detected paclobutrazol residue below permissible limit (0.4898–1.0005  $\mu\text{g/g}$ ) in the rhizosphere after two years of application. Jaradrattanapaiboon, et al. (2008) reported spatial difference of paclobutrazol residue in soils as they observed high concentration of PBZ residue in upper soil layer (0-5 cm) and low residue level in lower soil layer (10- 20 cm). They further reported that PBZ persisted for about 3-5 months. On the other hand, Narvaranant, et al. (2000) reported the persistence of PBZ residue up to 12 months Ochoa, et al. (2009) expressed the possibility of environmental contamination with the regular application of paclobutrazol in containerized oleander production due the leaching of PBZ into the nursery soil with the irrigation water. The adsorption and leaching of the residues is dependent upon the soil physical and chemical characteristics as well as environmental factors such as rainfall. Wu, et al. (2013) have reported that paclobutrazol was more persistent in greenhouse than in open field soil; leaching by rainfall being responsible for the difference in dissipation. Paclobutrazol is also known to leach in soil with high sand content.

## Conclusion

Paclobutrazol is a growth inhibitor and belong to triazol group. It inhibit the biosynthesis of GA3 at kaurene stage and it is most commonly used for the induction of flowering in off season, control tree vigour for HDP (canopy managment), increase fruit set and yield, improve fruit quality when applied to the soil. Studies aiming to adjust the amount of application dose of paclobutrazol to each crops will allow the formulation of recommendations for more efficient applications, which can not only provide quality fruit production throughout the year but also reduce the risk of residues in orchard soil, tree, fruit and environment. The cultar is most commonly used for the induction of flowering in off season, control tree vigour for HDP (canopy managment), increase fruit set and yield, improve fruit quality when applied to the soil but has the drawback of relatively high persistence in both soil and fruit in mango. Studies aiming to adjust the amount of application dose of cultar to each cultivar will allow the formulation of recommendations for more efficient applications, which can not only provide quality fruit production throughout the year but also reduce the risk of residues in the mango orchard soil, tree, fruit and environment.

## References

- Williams MW, Edgerton LJ, (1983) Vegetative growth control of apple and pear trees with ICI P-33 (paclobutrazol) a chemical analog of bayleton. *Acta Hort* 137:11-16. Link: <http://bit.ly/389b0nZ>
- Erez A (1984) Dwarfing peaches by pruning and by paclobutrazol. *Acta Hort* 146: 235-24. Link: <http://bit.ly/2Ktu5ZK>
- Aron Y, Monselise SP, Goren R, Costo J (1985) Chemical control of vegetative growth in citrus trees by paclobutrazol. *Hort Sci* 20: 96-98. Link: <http://bit.ly/3mjPLEU>
- Sarkar CB, Rahim MA (2012) Vegetative growth, harvesting time, yield and quality of mango (*Mangifera indica* L.) as influenced by soil drench application of paclobutrazol. *Bangladesh J Agri* 37: 335-348. Link: <https://bit.ly/3oZ5sTC>
- Upreti KK, Shivu Prasad SR, Reddy YTN, Rajeswara AN (2014) Paclobutrazol induced changes in carbohydrates and some associated enzymes during floral initiation in mango (*Mangifera indica* L.) cv. Totapuri. *Indian J Plant Physiol* 19: 317-323. Link: <https://bit.ly/37ptUYE>
- Anonymous (2017) NHB, Gurgaon, Haryana. Arteca, R. N. 1995. Plant growth substances: principles and applications. New York: Champman & Hall 332.
- Murti GSR, Upreti KK, Reddy YTN, Kurian RM (2000) Cytokinins and abscissic acid in xylem sap of mango cv. Alphonso grafted on rootstocks of contrasting vigour. *Indian J Hort* 57: 221-223. Link: <https://bit.ly/2KA036s>
- Patil PB, Rao MM, Srinivasan CN, Basarkar PW, Nalwadi VG (1992) Physiological and biochemical factors associated with fruit bud differentiation in Alphonso mango; V-total phenols and polyphenol oxidase. *Karnataka J Agric. Sci.* 5: 338-342.
- Protacio CM, Bugante, RD, Quinto J, Molinyawe G and Paelmo G (2000) Regulation of flowering in 'carabao' mango trees by paclobutrazol. *Philippine Journal of Crop Science* 25: 27-33. Link: <http://bit.ly/387TcJH>
- Blaikie SJ, Kulkarni VJ, Muller WJ (2004) Effects of morphactin and paclobutrazol flowering treatments on shoot and root phenology in mango cv. Kensington Pride. *Sci Hort* 101: 51-68. Link: <http://bit.ly/3r2Ndi5>
- Yeshitela T, Robbertse PJ, Stassen PJC (2004) Paclobutrazol suppressed vegetative growth and improved yield as well as fruit quality of 'Tommy Atkins' mango (*Mangifera indica*) in Ethiopia. *New Zealand J Crop Hort Sci* 32: 281-293. Link: <https://bit.ly/2LIAdOs>
- Nafeez M, Faqueer M, Ahmad S, Khan M, Jamil M, Aslam M (2010) Paclobutrazol soil drenching supresses vegetative growth, reduces malformation and increases production in mango. *Int J Fruit Sci* 10: 431-440. Link: <https://bit.ly/2WlwsXE>
- Burondkar MM, Rajan S, Upreti KK, Reddy YTN, Singh VK, et al. (2013) Advancing Alphonso mango harvest season in lateritic rockysoils of Konkan region through manipulation in time of paclobutrazol application. *Journal of Applied Horticulture* 15: 178–82. Link: <https://bit.ly/3agJiYY>
- Nartvaranant P, Subhadrabandhu S, Tongumpai P (2000) Practical aspect in producing off-season mango in Thailand. *Acta Hort* 509: 661-668. Link: <https://bit.ly/2Wi9tq9>
- Fonseca N, Castro NMT, Ledo CA (2004) Paclobutrazol on flowering and production of 'Tommy Atkins' mango. *Ciencia e Agrotecnologia* 28: 807- 814. Link: <http://bit.ly/3mxWLOn>
- Kim JK, Kim KY, Kim JB, Kim SB (1990) The effect of paclobutrazol on shoot growth, photosynthetic activity, leaf and fruit characteristics and flower bud formation in Fuji apples. *Res. Reports Rural Dev. Admins Hort* 32: 10-15. Link: <http://bit.ly/387j48J>
- Steffens GL, Jacobs FW, Engelhaupt ME (1993) Size, flowering and fruiting of maturing own-rooted 'Gala' apple trees treated with paclobutrazol sprays and trunk drenches. *Scientia Hort.* 56: 13-21. Link: <http://bit.ly/3ai6ICE>

18. Chaney WR (2005) Growth retardants: A promising tool for managing urban trees. Purdue Extension document FNR252-W. Link: <https://bit.ly/2Wj4W6z>
19. Kotur SC (2006) Effect of paclobutrazol on root activity of mango (*Mangifera indica*). *Ind J Agril Sci* 5: 75- 80. Link: <https://bit.ly/3mmPyRj>
20. Werner H (1993) Influence of paclobutrazol on growth and leaf nutrient content of mango. *Acta Horticulturae* 341: 255-259. Link: <http://bit.ly/3ntQDbp>
21. Singh NP, Malhi CS, Sharma RC (2005) Effect of plant bio regulators (PBRs) on flowering, fruit yield and quality in mango cv. Dashehari. *Horti J* 18: 10-12. Link: <https://bit.ly/3h0O0vq>
22. Upreti KK, Reddy YTN, Shivu Prasad SR, Bindu GV, Jayaram HL, et al. (2013) Hormonal changes in response to paclobutrazol induced early flowering in mango cv. Totapuri. *Sci Hort* 150: 414-418. Link: <http://bit.ly/2KAa03R>
23. Hedden P, Martin GC (1985) Inhibition of gibberellin biosynthesis by paclobutrazol in cell free homogenates of *Cucurbita maxima* endosperm and *Malus pumila* embryos. *J Plant Growth Reg* 4: 111-122. Link: <https://bit.ly/3nqIHZD>
24. Zhang ZH (1990) The effects of paclobutrazol on peach trees at the early fruiting stage. *Jiangsu Nongye Kexue* 1: 49-50.
25. Chen D, Shen D, Zailong L (1995) Effects of PP333 on early flowering of peach seedlings. *Acta Agric* 7: 20-23. Link: <http://bit.ly/3mthsHnd>
26. Allan P, George AP, Nissen RJ, Rasmussen TS, Morley Bunker MJ (1993) Effects of paclobutrazol on phenological cycling of low chill 'Flordaprince' peach in subtropical Australia. *Scientia Hort* 53: 73-84. Link: <https://bit.ly/3k7JOtr>
27. Biasi R, Costa G, Succi F (1989) Paclobutrazol and root zone water content influence peach seedling behaviour. *J Am Soc Hort Sci* 114: 923-926.
28. George AP, Nissen RJ, Rasmussen T (1993) Effects of postharvest topping, autumn cincturing, and paclobutrazol on growth, yield and fruit quality of the low chill nectarine cv. Sundowner in subtropical Australia. *Aust J Expt Agric*. 33: 353-362. Link: <http://bit.ly/34FnAdN>
29. Lever BG, Shearing SJ, Batch JJ (1982) PP333- A new broad spectrum growth retardant. *Proc. British Crop Protec. Conf. Weeds* 3-10.
30. Irving DE, Pallesen JC (1989) Growth retarding effects of paclobutrazol and RSWO 411 on Granny Smith and Fuji apple trees. *J Plant Growth Reg* 8: 199-204. Link: <http://bit.ly/2KBNkgg>
31. Stinchcombe GR, Copas E, Williams RR, Arnold G (1984) The effects of paclobutrazol and daminozide on the growth and yield of cider apple trees. *J Hort Sci* 59: 323- 327. Link: <https://bit.ly/3murLio>
32. El-Khoreiby AM, Unrath CR, Lehman LJ (1990) Paclobutrazol spray timing influences apple tree growth. *Hort Sci* 25: 310-312.
33. Greene DW (1986) Effect of paclobutrazol and analogs on growth, yield, fruit quality and storage potential of Delicious apples. *J Amer Soc Hort Sci* 111: 328- 332. Link: <http://bit.ly/2Wt8SBP>
34. Mavrodiev SG, Manolov PB (1989) A practical approach to use paclobutrazol (Cultar) on apples in Bulgaria. *Acta Hort* 239: 273-276 Link: <http://bit.ly/3gZV0Zo>
35. Webster AD, Quinlan JD, Richardson PJ (1986) The influence of paclobutrazol on growth and cropping of sweet cherry cultivar. *J Hort Sci* 61: 471-478. Link: <https://bit.ly/3myjNEW>
36. Arzani K, Bahaderi K, Piri S (2009) Paclobutrazol reduces vegetative growth and enhances flowering and fruiting of mature. *JH Link*
37. Brar JS (2010) Influence of paclobutrazol and ethephon on vegetative growth of guava (*Psidium guajava* L.) plants at different spacing. *Notulae Scientia Biologicae*. 2: 110-113. Link: <https://bit.ly/3gVdBWw>
38. Sharma MK, Joolka NK (2002a) Growth and water relations of Nonpareil almond as influenced by growth regulators. *The Hort. Journal* 15: 83-86.
39. Sharma MK, Joolka NK (2011) Influence of triacontanol and paclobutrazol on growth and leaf nutrient status of Non Pareil almond under different soil moisture regimes. *Indian Journal of Horticulture*. 68: 180-183. Link: <https://bit.ly/3gZqrTz>
40. Mir MM, Baba JA, Umar I, Rather GH, Rehman MU, et al. (2015) Effect of soil applied paclobutrazol on vegetative growth and quality attributes of apricot (*Prunus armeniaca* L.). *Green Farming Int. J.* 6: 813-813. Link: <https://bit.ly/3mv6c19>
41. Greene DW (1982) Effect of PP333 and its analogs on vegetative growth and fruit quality of Delicious apple. *Hort. Sci.* 17: 474.
42. Quinlan JD, Richardson PJ (1984) Effect of paclobutrazol (PP333) on apple shoots growth. *Acta Hort* 146:105-111. Link: <http://bit.ly/38dPfmR>
43. Khurshid T, Mcneil DL, Trought MCT (1997) Effect of foliar-applied gibberellins and soil-applied paclobutrazol on reproductive and vegetative growth of 'Braeburn' apple trees growing under a high-density planting system. *New Zealand Journal of Crop and Horticultural Science* 25. Link: <https://bit.ly/2WE3CM5>
44. Quinlan JD, Webster AD (1982) Effect of the growth retardant PP333 on growth of plums and cherries. *Int. Society Hort. Sci.* 1: 1071.
45. Klinac DJ, Rohitha H, Pevreal JC (1991) Effects of cultar (paclobutrazol) on vegetative growth and fruit production by Nashi (*Pyrus seratina* Rehd.). *New Zealand J Crop Hortl. Sci.* 19: 229-235. Link: <https://bit.ly/38bRd7q>
46. Young PS (1983) Peach growth response from PP333 (paclobutrazol). In: proceedings of tenth annual meeting plant growth regulator soc of america. East lansing, USA. PGR Society America, 192-194. Link:
47. Kulkarni VJ (1988) Chemical control of tree vigour and the promotion of flowering and fruiting in mango (*Mangifera indica* L.) using paclobutrazol. *J Hort. Sci.* 63: 557- 566. Link: <https://bit.ly/3aua6F4>
48. Tandell YN, Patel NL (2011) Effect of chemicals on growth, yield and economics of mango (*Mangifera indica* L.). *Karnataka J Agric Sci* 24: 362-365. Link: <https://bit.ly/3oWj95M>
49. Gonzalez A, Blaikie SJ (2003) Seasonal variation in carbon assimilation in mango (cv. Kensington Pride): Effect of flowering treatments. *Australian J Agri Res* 54: 309-321. Link: <https://bit.ly/2J3d08l>
50. Curry EA, Williams MW (1983) Promalin or GA3 increase pedicel and fruit length and leaf size of Delicious apples treated with paclobutrazol. *Hort. Sci.* 18: 214-215. Link:
51. Swietlik D, Miller SS (1985) The effect of paclobutrazol on mineral nutrition of apple seedlings. *J Plant Nutr* 8: 369-382. Link: <https://bit.ly/3nrqGD6>
52. Chogtu SK (1986) Effect of spacing and paclobutrazol (PP333) on photosynthetic efficiency, growth and yield of Royal Delicious apples. M.Sc. Thesis, Dr. Y.S. Parmar University of Horticulture and Forestry, Nauni, Solan, India. Link: <https://bit.ly/3p4UTP7>
53. Abod SA, Webster AD (1991) The influence of foliar sprays of tetcyclasis or paclobutrazol on the growth and water use of transplanted *Malus*, *Tilia* and *Bitula* stocks. *J Hort. Sci* 66: 85-94. Link: <https://bit.ly/3k7JdYJ>
54. Bhatia SK (1992) Effect of paclobutrazol, daminozide and rootstocks on the growth and chemical composition of apple plant. Ph.D. Thesis, Dr. Y.S. Parmar University of Horticulture and Forestry, Nauni, Solan, India.
55. Lenz F (1984) PP333- an interesting growth retardant. *Erweb.* 26:c244.
56. Steffens GL, Wang SY, Byun JK (1984) Effects of altering gibberellin availability on growth and chemical composition of apple seedlings. *Proc. 11th Annual Meeting, Pl. Growth Reg. Soc. America.* Link:
57. Lehman LJ, Young E, Unrath CR (1990) Apple tree vigour influences flowering and dry weight after paclobutrazol application. *Hort Sci.* 25: 933-935. Link: <http://bit.ly/3amQH98>
58. El-Hodairi MH, Canham AE, Buckley WR (1988) The effects of paclobutrazol

- on growth and the movement of <sup>14</sup>C-labelled assimilates in Red Delicious apple seedlings. *J Hort Sci* 63: 575-581. Link: <http://bit.ly/34j1zRu>
59. Huang WO, Shen T, Han ZH, Liu S (1995) Influence of paclobutrazol on photosynthesis rate and dry matter partitioning in the apple tree. *J Plant Nutr* 18: 901-910. Link: <https://bit.ly/3oXZfYa>
60. Curry EA, Williams MW (1990) Influence of rootstock on response of Delicious and Golden Delicious trees treated with paclobutrazol. *Fruit Var J* 44: 172-176. Link: <http://bit.ly/3awxL8f>
61. Swietlik D, Miller SS (1984) The effect of paclobutrazol on the growth and uptake of Ca and K by apple seedlings. *J Plant Nutr* 7: 1555-1556. Link: <https://bit.ly/2K5Yn58>
62. Zeller JK, Larson FE, Higgins SS, Raese JT, Fellman JK (1991b) Rootstock effects on responses of potted 'Smoothie Golden Delicious' apple to soil-applied triazole growth inhibitors. II. Mineral nutrition and carbohydrate status. *Scientia Hort* 46: 75-88. Link: <http://bit.ly/2KyQ4P6>
63. Singh RN (1958) Studies in the differentiation and development of fruit bud in mango. II Morphological and histological changes. *Hort Adv.* 2: 37. Link: <https://bit.ly/3r2wWJX>
64. Shinde AK, G.M Waghmare, RG, Wagh RG and M.M Burondkar (2000) Effect of dose and time of paclobutrazol application on flowering and yield of mango. *Indian J Plant Physiol* 5: 82-84. Link: <https://bit.ly/34I2a5k>
65. Chusri O, Kozai N, Ogata T, Higuchi H and Yonemoto Y (2008) Application of paclobutrazol for flowering and fruit production of 'Irwin' mango (*Mangifera indica* L.) in Okinawa. *Tropical Agriculture and Development* 52: 69-73. Link: <http://bit.ly/2WnCfFH>
66. Singh Z (2000) Effects of (2RS, 3RS) paclobutrazol on tree vigour, flowering, fruit set and yield in mango. *Acta Hort.* 525: 459-462.
67. Cardoso MGS, Jose ARS, Viana AES, Matsumoto SN, Reboucas TNH (2007) Flowering and production of mango fruits (*Mangifera indica* L) cv Rosa promoted by different paclobutrazol doses. Link: <http://bit.ly/3nsbvzZ>
68. Reddy YTN, Kurian RM (2008) Cumulative and residual effects of paclobutrazol on growth, yield and fruit quality of 'Alphonso' mango *Journal of Horticultural Sciences* 3: 119-122. Link: <http://bit.ly/3msdWkC>
69. Singh S, Singh AK (2006) Regulation of shoot growth and flowering in mango cv Gulab Khas by paclobutrazol. *Annals of Agricultural Research* 27: 4-8.
70. Mouco, MAdC, Albuquerque, JAS (2005) Paclobutrazol effect at two mango production cycles *Bragantia*, 64: 219-225. Link: <https://bit.ly/3r2vN53>
71. Bagel BS, Tiwari R, Gupta N (2004) Effect of cultar and NAA on flowering and fruiting of mango (*Mangifera indica* L) cv Langra. *South Ind Hort* 52: 302-304.
72. Elfying DC, Lougheed EC, Chu CL, Cline RA (1990) Effects of diaminozide, paclobutrazol and uniconazole treatment on 'McIntosh' apples at harvest and following storage. *J American Society Hortl Sci* 115: 750-756. Link: <https://bit.ly/3r6p5uL>
73. Stan S, Burloi N, Popescu I, Fenesanu N, Cotorobia M (1989) Performance of paclobutrazol (cultar) in controlling vegetative growth and cropping of store fruits. *Acta Hort* 239: 221-228. Link: <http://bit.ly/3nsXnq4>
74. Wolstenholme BN, Whiley AW, Saranah JB (1990) Manipulating vegetative, reproductive growth in avocado (*Persea americana* Mill) with paclobutrazol foliar sprays. *Scientia Hort* 41: 315-327. Link: <http://bit.ly/38dW0Ym>
75. Jindal KK, Chadel JS (1996) Effect of triacntanol and paclobutrazol on fruit set, growth and quality of *Prunus saliciana*. *Indian J Hort* 53: 262-268. Link: <https://bit.ly/3893IAE>
76. Ratna J, Bist LD (1997) Effect of cultar on growth, yield and fruit quality in grown up Gola pear trees. *Indian J Hort* 54:111-115. Link: <https://bit.ly/2Wn1VCj>
77. Arzani K, Wood DES, Lawes GS (2000) Influence of first season application of paclobutrazol, root- pruning and regulated deficit irrigation on second season flowering and fruiting of mature Sundrop apricot trees. *Acta Hort* 516: 75-82. Link: <https://bit.ly/34mMdlT>
78. Abdollahi M, Eshghi S, Tafazoli E (2010) Interaction of paclobutrazol, boron and zinc on vegetative growth, yield, and fruit quality of strawberry (*Fragaria × Ananassa Duch* cv Selva). *J Biol Environ Sci* 4: 67-75. Link: <https://bit.ly/3n0lwSS>
79. Fuentes AM, Mesejo C, Fambuena NM, Reig C, Mas MCG (2013) Fruit load restricts the flowering promotion effect of paclobutrazol in alternate bearing citrus spp. *Scientia Horticulturae* 151: 122-127. Link: <http://bit.ly/2KB4Btj>
80. Lina DP, Protacio CM (2015) Phenology of jackfruit (*Artocarpus heterophyllus* lam) and effects of paclobutrazol on off-season fruit production *Acta Hort* 1088: 495-500. Link: <https://bit.ly/3mrvqQn>
81. Singh VK, Singh A (2003) Effect of paclobutrazol on regularity of bearing in mango (*Mangifera indica* L). *Physiol Mol Biol Pl* 9: 239-248. Link: <https://bit.ly/2WraR9G>
82. Abdel RAOS, Osman ME, Fritz KB (2011) Effect of paclobutrazol (PBZ) on floral induction and associated hormonal and metabolic changes of biennially bearing mango (*Mangifera indica* L) cultivars during off year. *ARPN J Agri Bio Sci* 6: 55-67. Link: <https://bit.ly/32kZcwY>
83. Jogande AO, Chowdhary KG (2001) Seasonal changes in abscissic acid content and its role in flowering in mango (*Mangifera indica* L). *Orissa J Hort* 2001 29: 46-49.
84. Karki KB, Dhakal DD (2003) Effect of paclobutrazol on off-year induction of flowers in mango. *J Inst Agri Anim Sci* 24: 51-57. Link: <https://bit.ly/2LJVszr>
85. Singh DB, Ranganath HR (2006) Induction of regular and early fruiting in mango by paclobutrazol under tropical humid climate. *Indian J Hort* 63: 248-250. Link: <https://bit.ly/2Wlshob>
86. Hussien S, Kuswanto Ashari S, Basuki N (2012) Induction of flowering and yield of mango hybrids using paclobutrazol. *J Agril Food Tech* 2: 153-158. Link: <http://bit.ly/3p6gwyB>
87. Reddy YTN, Upreti KK, Prasad SR (2014) Response of paclobutrazol treatment on flowering, fruit maturity, yield and quality of mango (*Mangifera indica* L) cv Totapuri. *Indian J Agic Sci* 84: 1231-1235. Link: <https://bit.ly/2IXxTIH>
88. Vijaylakshmi D, Srinivasan, PL (2000) Improving the quality attributes of „off” year Alphonso mango through chemicals and growth regulators. *The Orissa J Hort* 28: 31-33. Link: <http://bit.ly/34hz8U8>
89. Hoda MN, Sanjay S, Jayant S (2001) Effect of cultar on flowering, fruiting and fruit quality of mango cv Langra. *Indian J Hort* 58: 224-227. Link: <https://bit.ly/3gYQ1ly>
90. Burondkar MM, Gunjate RT (1993) Control of vegetative growth and induction of regular and early cropping in 'Alphonso' mango with paclobutrazol. *Acta Hort* 341: 206-215. Link: <http://bit.ly/3nyxIMQ>
91. Kurian RM, Iyer CPA (1993) Stem anatomical characteristics in relation to tree vigour in mango (*Mangifera indica* L). *Sci Hort* 50: 245-253. Link: <http://bit.ly/3nw0OMw>
92. Suryanarayana V (1987) Seasonal changes in sugars, starch, nitrogen and C: N ratio in relation to flowering in mango. *Plant Biochem J* 5: 108-117. Link: <http://bit.ly/3nu5NNM>
93. Pongomboon W, Subhadrabandhu S, Stephenson RA (1997) Some aspects of the ecophysiology of flowering intensity of mango (*Mangifera indica* L) cv Nam Dok Mai in a semi-tropical monsoon Asian climate. *Sci Hort* 70: 45-56. Link: <http://bit.ly/3gWvTXi>
94. Ito A, Hayama H, Kashimura Y (2004) Possible role of sugar concentration and its metabolism in the regulation of flower bud formation in Japanese pear (*Pyrus pyrifolia*). *Acta Hort* 636: 365-373. Link: <http://bit.ly/3amNpTk>

95. Jyothi MH, Ravindra S, Suresh E, Soppin R, Ekbote S (2000) Biochemical changes in the leaf of bearing and nonbearing trees of some mango (*Mangifera indica* L) varieties/hybrids. *Adv Agri Res India* 10: 17-23. Link:
96. Palanichamy V, Reddy NN, Babu S, Selvaraj E Aranganathan, Mitra B (2012) Determination of time period of fruit bud differentiation and the associated histological and biochemical changes in mango hybrids. *Res JPharm Bio Chem Sci* 31: 271-289. Link: <https://bit.ly/2Wkr9RY>
97. Subhadrabandhu S, Tongumpai P, Kesta S, Suppkitjarak N (1997) Study of paclobutrazol on mango (*Mangifera indica*) cv Khiew Sawoey II Effect on total nonstructural carbohydrates, reducing sugars and total nitrogen contents in terminal shoots. *Thai J Agric Sci* 30: 269-282. Link: <http://bit.ly/3gTzxxY>
98. Rakshe MV, Nigade PM (2013) Pre and post flowering physiological behaviour of Alphonso mango leaves as influenced by stem girdling, root pruning and chemicals. *Bioinfolet* 10(3B): 979-982. Link: <https://bit.ly/2Kz1mCZ>
99. Phadung T, Krisanapook K, Phavaphutanon L (2011) Paclobutrazol, water stress and nitrogen induced flowering in 'Khao Nam Phueng' pummelo Kasetsart. *J Nat Sci* 45: 189-200. Link: <https://bit.ly/3aea0l4>
100. Davenport TL (2007) Reproductive physiology In: Litz RE (ed) *The Mango* Wallingford, CAB International; 97-169.
101. Abdel RAOS, Elamin OM, Bangerth FK (2008) Effects of paclobutrazol on floral induction and correlated phytohormonal changes in grafted seedlings of different mango (*Mangifera indica* L) cultivars Sudan *J Agri Res* 11: 111-120. Link: <https://bit.ly/3mXphJZ>
102. Shivu SR, Reddy YTN, Upreti KK, Rajeshwara AN (2014) studies on changes in carbohydrate metabolism in regular bearing and "off" season bearing cultivars of mango (*Mangifera indica* L) during flowering. *Int J Sci*; 52:148-152. Link: <https://bit.ly/2Wpoal0>
103. Murti GSR, Upreti KK, Kurian RM, Reddy YTN (2001) Paclobutrazol modifies tree vigour and flowering in mango cv Alphonso. *Indian J Plant Physiol* 6: 355-360. Link: <http://bit.ly/2WlpRpB>
104. Murti GSR, Upreti KK (2003) Endogenous hormones and phenols in rootstock seedlings of mango cultivars and their relationship with seedlings of mango cultivars and their relationship with seedling vigour. *Europ J Hort Sci* 68: 2-7. Link: <https://bit.ly/3gWuqQM>
105. Kumar M, Ponnuswami V, Jeya Kumar, Saraswathy S (2014) Influence of season affecting flowering and physiological parameters in mango. *Sci Res Essays* 9: 11-16. Link: <https://bit.ly/3qYlv4O>
106. Martin GC, Yoshikawa F, James HL (1987) Effect of soil application of paclobutrazol on growth, pruning time, flowering, yield and quality of Flavorcrest peach. *J American society for Hortl Sci* 112: 915-921.
107. Singh VK, Saini JP (2001) Regulation of flowering and fruiting in mango (*Mangifera indica* L) with paclobutrazol *Indian Society plant physiology* 61-68.
108. Saxena P, Singh VK, Pathak N (2013) Antioxidative enzymes and biochemical changes in paclobutrazol induced flowering in mango (*Mangifera indica*) cultivars. *J Envi Bio* 35: 1061-1066 Link: <http://bit.ly/3mnrYDX>
109. Bamini T, Parphiban S, Manivanan MI (2009) Effect of pruning and paclobutrazol on yield and quality of mango (*Mangifera indica* var Neelum). *Asian Sci* 4: 65-68 Link: <https://bit.ly/2WnfvZx>
110. Auxilia J, Sathiamoorthy S, Jeyakumar P, Kumar N, Balamohan TN (2010) Effect of paclobutrazol (PP333) on yield and quality of fruit and latex of Papaya var CO-2. *Acta Hort* 851: 413-418 Link: <https://bit.ly/2KvCj3J>
111. Jain MC, Dashora VX (2011) Effect of plant growth regulators on physico-chemical characters and yield of guava cv Sardar under high density planting system. *Indian J Hort* 68: 259-261. Link: <https://bit.ly/3r6FBLm>
112. EFSA 2006, (2018) Paclobutrazol Draft Assessment Report (DAR) Report and proposed decision of the United Kingdom made to the European Commission Available *Int. J. Curr. Microbiol. App Sci* 7: 1552-1562- 1561.
113. Benjawan C, Chutichudat P, Boontiang K, Chanaboon T (2006) Effect of chemical paclobutrazol on fruit development, quality and fruit yield of Kaew mango (*Mangifera indica* L) in Northeast Thailand. *Pakistan Journal of Biological Sciences* 9: 717-722. Link: <https://bit.ly/34k5KwH>
114. Anbu S, Parthiban S, Rajangam J and Thangaraj T (2001) Induction of off season flowering in mango (*Mangifera indica* L) using paclobutrazol *South Ind Hort*, 49: 384-385.
115. Hauser C, Kwiatkowski J, Rademacher W, Gossman K (1990) Regulation of endogenous abscisic acid levels and transportation in oilseed rape by plant growth retardants *J PI Physiol* 137: 201-207.
116. Kishore K, Singh HS, Kurian RM (2015) Paclobutrazol use in perennial fruit crops and its residual effects: A review. *Indian J Agril Sci* 85: 863-872. Link: <https://bit.ly/3amL8rg>
117. Aguirre R, Blanco A (1992) Patterns of histological differentiation induced by paclobutrazol and GA3 in peach shoots. *Acta Hort* 315: 7-12. Link: <https://bit.ly/32eD1bq>
118. Steffens GL, Wang SY, Faust M, Byun JK (1985) Growth, carbohydrate and mineral element status of shoot and spur leaves and fruit of Spartan apple trees treated with paclobutrazol. *J Am Soc Hort Sci.* 110: 850- 855. Link: <http://bit.ly/3oYbdkz>
119. Wieland WF, Wample RL (1985) Effects of paclobutrazol on growth, photosynthesis and carbohydrate content of Delicious apples. *Scientia Hort* 26: 139-147. Link: <http://bit.ly/3gX7bpR>
120. Curry EA (1988) Chemical control of vegetative growth of deciduous fruit trees with paclobutrazol and RSWO 411. *Hort Sci* 23:70-473.
121. Bonomo R, Neri D, Sansavini S (1989) Dry matter and nutrient partitioning, seasonal water consumption and gas exchange of paclobutrazol growth-controlled apple trees. *Acta Hort* 239: 159-166. Link: <http://bit.ly/2WqpKJs>
122. Sharma MK, Joolka NK (2003) Bioregulators mediated biochemical changes in Nonpareil almond under soil water stress Haryana. *J Hort Sci* 32: 182- 184.
123. Rieger M, Scalabrelli G (1990) Paclobutrazol, root growth, hydraulic conductivity and nutrient uptake of 'Nemaguard' peach. *Hort Sci* 25: 95-98. Link: <http://bit.ly/37oRcOe>
124. Allan P, George AP, Nissen RJ, Rasmussen TS (1995) Phenology and management practices of 'Flordaprince' peach. *Acta Hort* 409: 19-24. Link: <http://bit.ly/2KtAnsm>
125. Huett DO, George AP, Slack JM, Morris SC (1997) Diagnostic leaf nutrient standards for low-chill peaches in subtropical Australia *Aust. J Exp Agric* 37: 119-126. Link: <http://bit.ly/2WjV03>
126. Gracia DA (2014) Vegetative and reproductive development of 'Ataulfo' mango under pruning and paclobutrazol management. *J Agril Sci Tecnol* 16: 385-393. Link: <https://bit.ly/34kUcZY>
127. Ram, RB, Vismil, RS and Lal, B (2005) Effect of pruning severities and paclobutrazol on bearing behaviour of rejuvenated mango trees cv Dashehari. *New Agriculturist* 16: 1-9. Link: <http://bit.ly/3oZU9L8>
128. Anonymous (2010) *Agricultural Processed Food and Export Development Authority (APEDA)*, New Delhi.
129. Antognozzi E, Preziosi P, Standardi A (1987) Effects of paclobutrazol (PP333) on the vegetative activity and the course of photosynthesis in olive trees in the nursery *Annali della facolta di Agraria, Univ degli Studi di Perugia* 41: 827-838
130. Antognozzi E, Romani F (1989) Effects of paclobutrazol (PP333) and of pruning on the vegetative and productive activities of cv Golden Delicious apple trees. *Annali della facolta di Agraria, Univ degli studi di Perugia*; 41: 281-293. Link: <https://bit.ly/34qnnLb>

131. Hale' and 'Red Skin' peach trees *Horticulture. Environment and Biotechnology*; 50:84-93
132. Davenport TL (2008) Reproductive physiology In: Litz RE (ed), *The Mango, Botany, Production and Uses 2 nd Edition* CAB International.
133. Wallingford Oxon (2000) In press Davenport, TL, Processes influencing floral initiation and bloom: the role of phytohormones in a conceptual flowering model. *Hort Tech* 10: 733- 739
134. Davenport TL (2003) Management of flowering in three tropical and subtropical fruit tree species *Hort Sci*, 38: 1331–1335. Link: <https://bit.ly/34qnnLb>
135. Davenport TL (2007) Reproductive physiology of mango. *Braz J Plant Physio* 19: 363–376. Link: <http://bit.ly/38cFHsd>
136. Davenport TL (2009) Reproductive physiology In: Litz, RE (Ed), *The Mango: Botany Production and Uses*, 2nd edition CAB International, Wallingford, UK 97–169.
137. Davis TD (2000) Influence of triazole growth retardants on adventitious root formation in bean hypocotyl cutting *Proc Plant Growth Reg Soc Am* 13: 217-223
138. Deyton DE, Sams CE, Cummins JC (1991) Strawberry growth and photosynthetic responses to paclobutrazol *Hort Sci* 26: 1178-1180. Link: <https://bit.ly/3nv8cb9>
139. George AP, Nissen RJ, Campbell JA, Rasmussen T, Allan P, et al. (1995) Effects of paclobutrazol on growth and yield of low chill peaches in subtropical Australia. *Acta Hort*; 409:109-116. Link: <http://bit.ly/3mvCGs0>
140. Grochowska MJ, Hodum M, Mika A (2004) Improving productivity of four fruit species by growth regulators applied once in ultra-low doses to the collar. *Journal of Horticultural Science & Biotechnology* 79: 252- 259. Link: <https://bit.ly/3r7Jwrl>
141. Hu XS, Ding SY (1993) Effects of paclobutrazol on postharvest physiology and patterns of respiration and ethylene production of peach fruits. *Acta Agric* 19: 53-60. Link: <http://bit.ly/3gy9Mj8>
142. Lichou J, Devlavyay V, Garcin A, Vidaud J, Jay M (1988) Application of paclobutrazol to stone fruit trees Effects on vegetative growth and fruiting CTIFL-INRA. 215-217.
143. Pandey SN, Dinesh MR (2010) *Mango*. Indian Council of Agricultural Research, New Delhi 30- 97. Link: <http://bit.ly/2J4aQ8S>
144. Rajkumar M, Reddy YN, Chandrasekhar R, Srihari D (2007) Studies on the effect of foliar application of chemicals and growth regulators on the fruit yield of unpruned mango (*Mangifera indica* L) trees cv Baneshan The Orissa *J Hort* 3: 44-47.
145. Rath S, Rajput CBS (2000) Effect of B naphthoxy acetic acid and gibberellic acid on chemical composition of mango fruits. *Orissa J Agric Res* 3: 155-156. Link: <http://bit.ly/2K5t7DI>
146. Sharma MK, Joolka NK (2002) Effect of bioregulators and rootstocks on growth, photosynthetic efficiency and water relations of Nonpareil almond *Annals of Agricultural Research* 23: 740-743.
147. Sharma RR, Singh R (2006) Pruning intensity modifies canopy microclimate, and influences sex ratio, malformation incidence and development of fruited panicles in „Amrapali” mango (*Mangifera indica* L). *Scientia Horticulturae* 109: 118–122. Link: <https://bit.ly/38coFdT>
148. Singh Z, Dhillon BS (1992) Effect of paclobutrazol on floral malformation, yield and quality of mango (*Mangifera indica* L). *Acta Hort* 296: 51-53. Link: <http://bit.ly/3anpdQE>
149. Singh DB, Ranganath HR (2006) Induction of regular and early fruiting in mango by paclobutrazol under tropical humid climate. *Ind J Hort* 63: 248-249. Link: <https://bit.ly/2Wlshob>
150. Singh S, Singh AK (2006) Regulation of shoot growth and flowering in mango cv Gulab Khas by paclobutrazol. *Annals of Agricultural Research* 27: 4-8.
151. Singh Z, Muller W, Polesny F (ed), Verheyden C (ed), Webster AD (2000) Effect of (2RS, 3RS) paclobutrazol on tree vigour, flowering, fruit set and yield in mango Proceedings of the international conference on integrated fruit production, Leuven, Belgium, 27 July-1 August 1998 *Acta-Horticulturae*, 525: 459-462.
152. Srivastava RP, Kumar S (2009) *Fruit and vegetable preservation: Principles and Practices* 360.
153. Reddy SA, Rajkumar M, Padmaja A, Rao PV (2010) Effect of plant growth regulators on flowering and yield of watermelon (*Citrullus lanatus* T). *J Hort Sci and Ornamental plants* 2: 19-23.
154. Whiley, AW, Wolstenholme BN, Saranah JB (1988) Canopy management toward 2000 “Avocados Toward 2000” Australian Avocado Bicentennial conference. At Caloundra, Australia 44-50.
155. Zora S, Singh Z, Muller W, Polesny F, Verheyden C, et al (2000) Effect of (2RS, 3RS) paclobutrazol on tree vigour, flowering, fruit set and yield in mango *Acta Hort* 525: 459-462. Link: <https://bit.ly/3l51mli>